

Growth Model Runoff II

Greg Johnson

Weyerhaeuser Company

December 15, 2005, Corrected September 17, 2006

Introduction

GROWTH MODEL RUNOFF II is a follow-up to our first Runoff conducted in 2002. We are interested in looking at the performance of commonly available Pacific Northwest Douglas-fir growth models in projecting stands with thinning and fertilization treatments. We are not looking for “truth” (although that would be great) – just to understand how each model simulates treatment effects.

Introduction

In **GMRO I** [Johnson (2002)] users were asked to run three stands through the growth model they used and report the results. The hypothesis under test was that there would be no significant user effects, but potentially large model effects.

We found that there were both **user** and **model** effects. The same model and version produced different results depending on the user. Model effects were sometimes large.

Introduction

GMRO II extends the experiment to examine how Douglas-fir growth models handle thinning and fertilization. In this experiment, user effects were minimized by asking each model's author to make the projections. The thought was to limit potentially misleading user effects from the main effect of the model.

Models Examined

The models examined are:

- DFSIM v1.4 (Marshall)
- FPS v6.44 (Arney)
- FVS (Smith-Mateja)
- SMC-ORGANON v8.0 (Hann)
- SPS 4.1h, circa 1999 (Hamlin)
- TASS vII (Goudie)

Models Examined

DFSIM

- **Model:** Douglas-Fir **SIM**ulator [Curtis, et. al. (1981)]
- **Version:** 1.4
- **Type** [Munro (1974)]: Whole Stand
- **Settings:**
 - Specified residual stand rather than specifying the residual basal area per acre (BA) and trees per acre (TPA).
 - Starting conditions specified the total age, TPA and BA.
 - The maximum Curtis relative density (RD) was allowed to increase above the default value of 70 up to 100.

Models Examined

FPS

- **Model:** Forest Projection System [Arney (2005)]
- **Version:** 6.44
- **Type:** Individual tree - distance dependent.
- **Settings:**
 - Region 12 - Western Washington.
 - Used composite tree list – did not assume each tree record was an individual tree.
 - Assumed a residual thinning clumpiness of 0.92
 - Site prep setting: 8Y75

Models Examined

FVS

- **Model:** Forest Vegetation Simulator [Donnelly (1997)]
- **Version:** Pacific Northwest Coast (PN) Variant
- **Type:** Individual tree - distance independent.
- **Settings:**
 - Location Run: Olympia
 - SDI_{max} of 520
 - BA_{max} of 300
 - This variant does not have keywords applicable to fertilizer in Oregon. Used keywords to increase basal area growth by 30% for 10 years and decreased mortality by 10% for 10 years. After 10 years growth and mortality defaulted to unmodified growth equations.

Models Examined

SMC-ORGANON

- **Model:** Stand Management Cooperative - ORGANON [Hann (2003)]
- **Version:** 8.0
- **Type:** Individual tree - distance independent.
- **Settings:**
 - No special settings reported.

Models Examined

SPS

- **Model:** Stand Projection System [Arney (1985)]
- **Version:** 4.1h
- **Type:** Individual tree - distance independent.
- **Settings:**
 - No special settings reported.

Models Examined

TASS

- **Model: Tree And Stand Simulator [Mitchell (1975)]**
- **Version: II**
- **Type:** Individual tree - distance dependent.
- **Settings:**
 - Expanded tree records to create a pseudo-fixed area plot (TASS requires a fixed-area plot).
 - Randomly thinned out a proportion of the replicated trees based on the probability of thinning shown in the “after” tree list.
 - Randomly added or removed a few trees from the expanded after-thinning tree list to ensure a target of 180 trees/ac (445/ha) was achieved.
 - Since TASS is spatially explicit, we established the unthinned trees to ensure they were well spaced, and then randomly added the thinned trees in between.
 - Created a unique dominant height growth curve that starts at the establishment height and passes through the assigned height at breast height age 50.
 - The fertilizer response model was developed using our large fertilizer-thinning experiment in coastal BC. The data and models indicate little or no response for sites above 35m (114') at BH age 50. Thus, the fertilization runs for Sites 1 and 2 were not done because TASS shows no response for these high sites.

Three stands were provided spanning a range of site qualities; young and high enough stocking to be thinned. The initial conditions were:

Stand	Site	TPA	BA	Dq	HT	HTLC	RD	Age_{bh}	Age_{total}
1	120	608	146.9	6.7	58.2	31.6	56.9	22	29
2	100	732	138.4	5.9	50.1	20.4	57.0	17	25
3	140	384	151.5	8.5	59.7	27.1	51.9	17	19

Stands were sub-sampled for height and crown ratio. For trees without field-measured heights and crowns, regressions specific to each stand were used to impute missing values.

The Problem

After Thinning

The thinnings were all to 180 trees per acre with a $\frac{d}{D}$ less than 1.0.
The after thinning stand statistics were:

Stand	Site	TPA	BA	Dq	HT	HTLC	RD
1	120	180	63.1	8.0	64.1	32.8	22.3
2	100	180	47.8	7.0	54.3	21.2	18.1
3	140	180	86.3	9.4	62.5	27.5	28.2

The Theory

Both thinning and fertilization have a long history of use in Pacific Northwest Douglas-fir silviculture. There is a rich resource of designed experiments and response-surface studies devoted to quantifying the effects of these treatments. Given this background, we would expect a clear understanding of these treatments.



The Theory

Thinning

Conventionally thinning is thought to concentrate growth on fewer stems per acre. Thus, while possibly not increasing total volume production, stem size is enhanced with treatment. Existing research is equivocal on how height growth is affected:


- A study at Wind River [Harrington and Reukema (1983)] found that pre-commercial thinning had a shock effect on height growth resulting in substantial height growth reductions.
- The Levels of Growing Stock Study (LOGS) [Marshall and Curtis (2002)] found stable height growth trends, with thinning treatments resulting in higher exhibited site index over time (possibly due to the chain-saw effect).



The Theory

Thinning

A few studies have shown an increase in total volume production with thinning. Some of the LOGS [Marshall and Curtis (2002)] light thinning treatments have now exceeded total volume production on the controls. Generally, the reduction in growing stock from thinning reduces total volume growth. This is equivalent to moving left along the Langsaeter curve [Langsaeter (1944)] prior to the plateau (regions I and II).



The Theory

Fertilization

Nitrogen fertilization of Douglas-fir has been studied for several decades, primarily through the Regional Forest Nutrition Research Program (RFNRP). The RFNRP found that Douglas-fir generally responded to additions of Nitrogen as urea. Many soils in the Pacific Northwest have sufficiently low available N for economic responses to be obtained. In a summary of RFNRP and other research [Miller, et. al. (1986)] documented the type and duration of response. Of the findings, several are important:

- Response duration was limited – the effect of N additions was not permanent.
- Response is inversely proportional to site quality as measured by site index (at least in unthinned stands).
- Absolute and relative response was greater in thinned stands.
- No mention of height growth response.



The Theory

Fertilization

Another RFNRP report [Stegemoeller and Chappell (1988)] documented 16 year results on fertilization of unthinned and thinned stands. Installations ranged between site index 114 and 118, and averaged $200 \frac{\text{ft}^2}{\text{acre}}$ basal area unthinned and $120 \frac{\text{ft}^2}{\text{acre}}$ thinned. Conclusions from the report:

- Response duration was limited, ranging from 8 years in thinned stands to 14 years in unthinned conditions.
- Basal area response averaged $10.8 \frac{\text{ft}^2}{\text{acre}}$ in unthinned stands and $7.4 \frac{\text{ft}^2}{\text{acre}}$ in thinned stands over a 16 year period.
- Volume response averaged $455 \frac{\text{ft}^3}{\text{acre}}$ in unthinned stands and $406 \frac{\text{ft}^3}{\text{acre}}$ in thinned stands over a 16 year period.
- No mention of height growth response.

Results

Definitions:

- Growth:

$$Growth_t = (X_t - X_0)$$

- Cumulative Fertilized Response:

$$Response_t = (Treated_{t,i} - Control_{t,i})$$

where t = time period, i = thinning treatment

- Cumulative Thinned Response:

$$Response_t = (Treated_t - Control_t) + (Control_0 - Treated_0)$$

Cubic Volume Growth after 20 Years

Site	Model	NN	TN	NF	TF
100	DFSIM	5398	4448	6244	5301
100	FPS v6.44 Western Washington	4364	2104	6858	3355
100	FVS PN OLY	4502	3157	4978	3558
100	ORGANON—SMC	4603	3285	4798	3495
100	SPS 4.1h	5104	3087	5229	3459
100	TASS	4859	3687	5959	4316
120	DFSIM	5520	4830	6026	5310
120	FPS v6.44 Western Washington	5084	2983	6950	4158
120	FVS PN OLY	3817	3277	4372	3739
120	ORGANON—SMC	5066	4126	5186	4268
120	SPS 4.1h	6164	3897	6738	4481
120	TASS	5959	5116		
140	DFSIM	10168	10237	10904	11084
140	FPS v6.44 Western Washington	9651	6729	11533	8246
140	FVS PN OLY	4904	4392	5523	4964
140	ORGANON—SMC	6793	5803	6878	5905
140	SPS 4.1h	8713	6919	9111	7761
140	TASS	8675	7931		

where: NN = No Treatments, TN = Thinned, NF = Fertilized, TF = Thinned and Fertilized.

Basal Area Growth after 20 Years


Site	Model	NN	TN	NF	TF
100	DFSIM	117	127	136	150
100	FPS v6.44 Western Washington	91	47	135	70
100	FVS PN OLY	126	92	138	105
100	ORGANON—SMC	90	88	94	94
100	SPS 4.1h	97	73	99	85
100	TASS	122	100	146	115
120	DFSIM	87	99	96	108
120	FPS v6.44 Western Washington	84	55	115	75
120	FVS PN OLY	99	83	111	94
120	ORGANON—SMC	77	88	79	91
120	SPS 4.1h	107	78	117	93
120	TASS	139	127		
140	DFSIM	187	229	202	249
140	FPS v6.44 Western Washington	172	125	200	149
140	FVS PN OLY	97	92	109	104
140	ORGANON—SMC	95	106	97	108
140	SPS 4.1h	147	136	156	159
140	TASS	195	185		


where: NN = No Treatments, TN = Thinned, NF = Fertilized, TF = Thinned and Fertilized.


Dominant Height Growth after 20 Years


Site	Model	NN	TN	NF	TF
100	DFSIM	38.0	38.0	40.6	40.6
100	FPS v6.44 Western Washington	30.1	30.6	42.3	42.7
100	FVS PN OLY	31.6	31.7	31.1	31.6
100	ORGANON—SMC	31.2	29.4	31.7	29.5
100	SPS 4.1h	33.9	34.9	32.9	35.6
100	TASS	35.1	34.8	38.7	38.4
120	DFSIM	41.7	41.7	43.6	43.6
120	FPS v6.44 Western Washington	33.3	34.6	43.7	45.1
120	FVS PN OLY	33.6	38.2	35.9	38.7
120	ORGANON—SMC	37.3	34.2	37.7	34.6
120	SPS 4.1h	38.6	39.7	39.4	40.7
120	TASS	37.1	38.1		
140	DFSIM	58.3	58.3	59.8	59.8
140	FPS v6.44 Western Washington	49.5	51.2	58.3	60.0
140	FVS PN OLY	46.8	49.0	48.7	49.8
140	ORGANON—SMC	48.1	45.8	48.1	45.9
140	SPS 4.1h	50.2	51.1	50.0	51.0
140	TASS	53.5	52.8		


where: NN = No Treatments, TN = Thinned, NF = Fertilized, TF = Thinned and Fertilized.






















































































Conclusions

The six models tested here show a wide range of responses to thinning, fertilization, and the combination of those treatments. No one model adhered to all of general research findings on these treatments. Users need to be careful when applying these models to silviculture investment analyses.

Of the models tested, TASS is the closest to a process-level model. It was well behaved, however, because of its base dataset, it was unable to estimate fertilization on medium to high PNW Douglas-fir sites.

Observations about Thinning

- Two models predict basal area response incremental to the control.
- One model predicts reductions in height growth with thinning.
- One model predicts “bonus wood” – total volume growth with thinning exceeds the control.
- Half the models have begun to accrete volume relative to the controls by the end of the 20 years.

Observations about Fertilization

- One model predicts very large increases in basal area and dominant height with fertilization.
- Most models show a peak response at 10 years from application.
- Three models are in the cubic volume ballpark suggested by the RFNRP report for sites between 100 and 120 but appear to have a high basal area growth response. Is height growth response being underestimated?
- Almost all models show decreased response with thinning.
- Several models show increased response to fertilization as site index increases.





Thanks to the following for making the projections:

- Jim Arney – FPS
- Jim Goudie – TASS
- Dave Hamlin – SPS
- David Hann – ORGANON
- David Marshall – DFSIM
- Erin Smith-Mateja – FVS

Literature Cited



Arney, J.D. 1985. *A modeling strategy for the growth projection of managed stands*. Can. J. For. Res. 15:511-518.



Arney, J.D. 2005. *Forest Projection System*. 2005. Forest Biometrics Institute
<http://www.forestbiometrics.com/>



Curtis, Robert O., Gary W. Clendenen, and Donald J. Demars. 1981. *A new stand simulator for coast Douglas-fir: DFSIM user's guide*. USDA For. Serv. Gen. Tech. Rep. PNW-128, 79 p. Pacific Northwest Forest and Range Experiment Station, Portland, Oregon.



Donnelly, D.M. 1987. *PACIFIC NORTHWEST COAST VARIANT of the Forest Vegetation Simulator*. WO-Forest Management Service Center, USDA-Forest Service Fort Collins, Colorado.



Hann, D.W., D.D. Marshall, and M.L. Hanus. 2003. *Equations for predicting height-to-crown-base, 5-year diameter-growth rate, 5-year height-growth rate, 5-year mortality rate, and maximum size-density trajectory for douglas-fir and western hemlock in the coastal region of the Pacific Northwest*. Forest Research Lab., Oregon State University, Corvallis, Oregon. Research Contribution 40. 83p.



Harrington, C.A., and D.L. Reukema. 1983. *Initial shock and long-term stand development following thinning in a Douglas-fir plantation*. Forest. Sci. 29:33-46.



Johnson, G.P. 2002. *Growth Model Users Group Growth Model Run-off*.
www.growthmodel.org/papers/modelrunoff.ppt.

Literature Cited



Langsaeter, A. 1944. *Om tynning i enaldret gran-og furuskog*. Referat: Produktionsundersuchungen von Fichtenwald. Medd. Det norske Skogforsoksvesen. VIII: 131-216.



Marshall, D.D. and R. O. Curtis. 2002. *Levels-of-Growing-Stock Cooperative Study in Douglas-fir: Report No. 15Hoskins: 19631998* Pacific Northwest Research Station Research Paper PNW-RP-537.



Miller, R.E, P. R. Barker, C.E. Peterson, and S.R. Webster. 1986. *Using Nitrogen fertilizers in management of coast Douglas-fir. I. Regional Trends of Response*. In: Using Nitrogen Fertilizer in Management of Coast Douglas-fir, RFRNP Report Number 7. College of Forest Resources, University of Washington.



Mitchell, K.J. 1975. *Dynamics and simulated yield of Douglas-fir*. For. Sci. Monogr. 17, 39 pp.



Munro, D.D. 1974. *Forest growth models: A prognosis*. In: Fries, J. (ed.) Growth Models for Tree and Stand Simulation. Royal College of Forestry, Department of Forest Yield Research, Stockholm, no. 30 in Research Notes, pp. 721.



Stegemoeller, K.A. and H.N. Chappell. 1988. *Growth response to single and multiple applications of N fertilizer over 16 years in unthinned and thinned Douglas-fir stands in the Pacific Northwest*. RFNRP Report Number 11. College of Forest Resources, University of Washington.